

DATA 1010
IN-CLASS EXERCISES
SAMUEL S. WATSON
05 NOVEMBER 2018

Problem 1

Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is a smooth function whose values on a grid of points $x_s = a : \epsilon : b$ are stored in a vector v . Write a Julia expression which approximates $\int_a^b f$.

Problem 2

Suppose that $f : [a, b] \rightarrow \mathbb{R}$ is a probability density function whose values on $x_s = a : \epsilon : b$ are stored in a vector v . Write a Julia expression which approximates $\mathbb{E}[X]$, where X is a random variable whose density is f .

Problem 3

Given a flower randomly selected from a field, let X_1 be its petal width in centimeters, X_2 its petal length in centimeters, and $Y \in \{R, G, B\}$ its color. Let

$$\begin{aligned} \boldsymbol{\mu}_R &= \begin{bmatrix} 9 \\ 5 \end{bmatrix} & \boldsymbol{\mu}_G &= \begin{bmatrix} 4 \\ 10 \end{bmatrix} & \boldsymbol{\mu}_B &= \begin{bmatrix} 7 \\ 9 \end{bmatrix} \\ A_R &= \begin{bmatrix} 1.5 & -1 \\ 0 & 1 \end{bmatrix} & A_G &= \begin{bmatrix} 0.5 & 0.25 \\ 0 & 0.5 \end{bmatrix} & A_B &= \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}. \end{aligned}$$

Suppose that the joint distribution of X_1, X_2 , and Y has the property that for any $A \subset \mathbb{R}^2$ and color $c \in \{R, G, B\}$, we have

$$\mathbb{P}(A \times \{c\}) = p_c \int_{\mathbb{R}^2} f_c(x_1, x_2) dx_1 dx_2,$$

where $(p_R, p_G, p_B) = (1/3, 1/6, 1/2)$ and f_c is the multivariate normal density with mean $\boldsymbol{\mu}_c$ and covariance matrix $A_c A_c'$.

Find the best predictor of Y given $(X_1, X_2) = (x_1, x_2)$ (using the 0-1 loss function), and find a way to estimate that predictor using the given samples.